ARTICLE · 2011

EMDR and Developmental Trauma

By Morit Heitzler

By Morit Heitzler. An examination of how EMDR — developed primarily for single-event PTSD — can and cannot be applied to developmental trauma, and what modifications to standard EMDR protocol are required when working with clients whose trauma is relational and cumulative rather than event-based.

KEY THEMES

— EMDR in single-event vs developmental trauma
— The limits of standard EMDR protocol
— Relational attunement as the primary variable
— Modifying EMDR for complex presentations
— Integration of EMDR with relational and somatic approaches

EMDR's Strengths and Limits

EMDR has established itself as one of the most evidence-based approaches to trauma treatment for single-event PTSD. Its effectiveness in helping clients process discrete traumatic memories — events that are stored in the nervous system in an unprocessed form, generating intrusive symptoms — is well-documented. The bilateral stimulation component, whatever its ultimate mechanism of action, seems to facilitate a form of memory processing that reduces the intensity of the traumatic response.

Where EMDR runs into difficulty is with developmental trauma — the cumulative relational injuries of childhood that do not take the form of discrete, identifiable events but are woven into the very fabric of the client's personality and relational patterns. For these clients, there is often no single 'trauma memory' to process; the trauma is not a memory but a way of being in the world.

Applying standard EMDR protocol to such clients without modification is at best ineffective and at worst actively destabilising — not because EMDR is the wrong tool, but because the therapeutic relationship which EMDR protocol assumes (a relatively stable, trusting, well-resourced client working with a technically competent therapist) is precisely what developmental trauma makes unavailable.

The Primacy of the Therapeutic Relationship

As soon as developmental trauma is involved, what really matters is the client's implicit and unconscious experience of the therapeutic relationship, regardless of the therapist's input. This is not a criticism of EMDR but a recognition that it was not designed with the most complex clients in mind.

The therapeutic relationship in developmental trauma work is not a warm background against which EMDR processing happens — it is itself the primary therapeutic instrument. The client's nervous system is regulated (or dysregulated) primarily by the quality of the relational field, not by the specific technique being employed within it.

This means that before EMDR can be effectively used with developmental trauma clients, a substantial period of relational work is required — establishing the therapeutic relationship as a reliable resource, developing the client's capacity for somatic self-awareness and dual attention, and building a working alliance that can survive the inevitable ruptures and repairs of complex trauma work.

Modifications Required

When EMDR is used with developmental trauma clients who have developed sufficient relational stability, several modifications to standard protocol are typically required. The pace of processing is typically slower; the interweaves needed more frequent; the somatic dimension of the processing more central.

Most importantly, the therapist needs to be willing to move outside the EMDR protocol when the relational field indicates that the protocol itself has become part of the enactment — when the structure of the bilateral stimulation and the therapist's direction of the client's attention is being experienced (unconsciously) as another version of the invasion and control that characterised the original trauma.

The integrated approach described in the companion articles involves using EMDR as one element within a broader relational and somatic framework — not as the primary treatment modality but as a tool to be used selectively, in response to the client's developing capacity, within a relationship that remains central throughout.

BIBLIOGRAPHY

— Shapiro, F. (1989) EMDR: Basic principles. Journal of Traumatic Stress

— Herman, J. (1992) Trauma and Recovery. New York: Basic Books

— Heitzler, M. (2009) Towards an integrative model of trauma therapy. In L. Hartley (ed.) Contemporary Body Psychotherapy. Routledge

— van der Kolk, B. (2014) The Body Keeps the Score. New York: Viking

— Lyons-Ruth, K. (1998) Implicit relational knowing. Infant Mental Health Journal 19

Published at soth.co.uk · © 2011 Morit Heitzler · All rights reserved